



EVERYDAY CHICK STARTER

MEDICATED Type C

ACTIVE DRUG INGREDIENT: Amprolium .0125%

FEEDING DIRECTIONS: As an aid in prevention of coccidiosis where immunity to coccidiosis is not desired. Feed to replacements up to 10 weeks of age.

WARNING: Use as a sole source of Amprolium.

SPECIAL CONSIDERATION: Do not use Amprolium in feeds containing bentonite.

The suggested feeding program is for use as a guide only. The animal's requirements may change due to breed, environment and management.

GUARANTEED ANALYSIS

| | | | |
|---------------|-------|-------|-------------|
| CRUDE PROTEIN | | (MIN) | 21.0% |
| LYSINE | | (MIN) | 1.00% |
| METHIONINE | | (MIN) | 0.35% |
| CRUDE FAT | | (MIN) | 3.50% |
| CRUDE FIBER | | (MAX) | 4.50% |
| CALCIUM | (MIN) | 1.10% | (MAX) 1.30% |
| PHOSPHORUS | | (MIN) | 0.50% |
| SALT | (MIN) | 0.20% | (MAX) 0.30% |
| SODIUM | (MIN) | 0.08% | (MAX) 0.15% |

INGREDIENTS

Ground Yellow Corn, Soybean Meal, Corn Gluten Meal, Wheat Middlings, Corn Distillers Dried Grains, Fish Meal, Fat-Vegetable Blend, Dehydrated Alfalfa Meal, Salt, Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Choline Chloride, Manganous Oxide, Zinc Oxide, Iron Carbonate, Niacin, Mineral Oil, Vitamin A, dl-Methionine, Sodium Selenite, Iron Sulfate, Zinc Sulfate, Pantothenic Acid, Vitamin B₁₂, Copper Sulfate, Riboflavin, Vitamin D, Vitamin E, Menadione Sodium Bisulfite (source of Vitamin K), Calcium Iodate, Copper Oxide, Cobalt Carbonate, Thiamine, Flavoring.

Ingredients as defined by AAFCO.

Do not store on ground, store in a cool, dry place.

Manufactured by:



148 Longmeadow Road, Taunton, MA 02780