



EVERYDAY CHICK STARTER MEDICATED Type C

ACTIVE DRUG INGREDIENT: Amprolium .0125%

INDICATIONS FOR USE: For the prevention of coccidiosis where immunity to coccidiosis is not desired.

FEEDING DIRECTIONS: Food continuously from day old until onset of production.

WARNING: Keep this and all drugs out of the reach of children.

The suggested feeding program is for use as a guide only. The animal's requirements may change due to breed, environment and management.

GUARANTEED ANALYSIS

CRUDE PROTEIN		(MIN)	21.0%
LYSINE		(MIN)	1.00%
METHIONINE		(MIN)	0.35%
CRUDE FAT		(MIN)	3.00%
CRUDE FIBER		(MAX)	4.50%
CALCIUM	(MIN)	1.00%	(MAX) 1.25%
PHOSPHORUS		(MIN)	0.50%
SALT	(MIN)	0.20%	(MAX) 0.30%
SODIUM	(MIN)	0.08%	(MAX) 0.15%

INGREDIENTS

Ground Yellow Corn, Soybean Meal, Wheat Middlings, Distillers Dried Grains w/ Solubles, Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Salt, dl-Methionine, Roughage Products, Vitamin E Supplement, Niacin Supplement, Calcium Pantothenate, Riboflavin Supplement, Biotin, Menadione Nicotinamide Bisulfite, Vitamin A Supplement, Vitamin D3 Supplement, Pyridoxine Hydrochloride, Thiamine Mononitrate, Mineral Oil, Folic Acid, Vitamin B₁₂ Supplement, Processed Grain By-Products, Manganous Oxide, Zinc Oxide, Ferrous Sulfate, Copper Sulfate, Natural Flavors, Sodium Selenite, Ethylenediamine Dihydriodide, Choline Chloride, l-Lysine, l-Threonine.


Ingredients as defined by AAFCO

Do not store on ground, store in a cool, dry place.

CAUTION

- Use as the sole source of amprolium.
- Do not change the litter while giving this feed unless absolutely necessary.
- If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis and follow the instructions of your veterinarian or poultry pathologist.
- Losses may result from intercurrent disease or other conditions affecting drug intake which can contribute to the virulence of coccidiosis under field conditions.
- In replacement flocks, the grower must expect that excessive exposure to one or more species may overwhelm the drug in some flocks and prompt treatment will be required.
- Do not use in feeds containing bentonite.

Manufactured by:



148 Longmeadow Road
Taunton, MA 02780
(508) 824-7292