



# EVERYDAY GOAT 16%

## STARTER/GROWER PELLETS

MEDICATED Type C - ACTIVE DRUG INGREDIENT: Decoquinatate 22.7 g/ton

### INDICATIONS FOR USE

With Decoquinatate and Yeast Culture. Also contains ammonium chloride to aid in prevention of kidney stones in young male goats.

**Goats:** For the prevention of coccidiosis in young goats caused by Eimeria christenseni and E. ninakohlyakimovae.

### GUARANTEED ANALYSIS

CRUDE PROTEIN	(MIN)	16.0%		
CRUDE FAT	(MIN)	3.50%		
CRUDE FIBER	(MAX)	10.0%		
ACID DETERGENT FIBER	(MAX)	15.0%		
CALCIUM	(MIN)	1.25%	(MAX)	1.50%
PHOSPHORUS	(MIN)	0.60%		
SALT	(MIN)	0.55%	(MAX)	0.65%
COPPER	(MIN)	15ppm	(MAX)	25ppm
SELENIUM	(MIN)	0.50ppm		
VITAMIN A	(MIN)	7000 IU/lb		

### INGREDIENTS

Wheat Middlings, Soybean Hulls, Ground Yellow Corn Meal, Corn Distillers Dried Grains w/ Solubles, Cane Molasses, Soybean Meal, Calcium Carbonate, Ammonium Chloride, Salt, Yeast Culture, dl-Alpha-Tocopheryl Acetate (Source of Vitamin E), Manganous Oxide, Zinc Oxide, Ferrous Sulfate, Biotin Supplement, Choline Chloride, Mineral Oil, Thiamine Mononitrate, Sodium Selenite, Riboflavin Supplement, Niacin Supplement, Vitamin A Acetate, Folic Acid Supplement, Calcium Pantothenate, Vitamin D<sub>3</sub> Supplement, Pyridoxine Hydrochloride, Vitamin B<sub>12</sub> Supplement, Calcium Iodate, Menadione Sodium Bisulfite Complex, Cobalt Carbonate, Manganous Sulfate, Zinc Sulfate, Selenium Yeast, Basic Copper Chloride, Natural Flavors, Ethylenediamine Dihydriodide, Fat-Vegetable Blend, Potassium Sulfate, Magnesium Sulfate, Monocalcium Phosphate, Dicalcium Phosphate, dl-Methionine.

**Ingredients as defined by AAFCO. Do not store on ground, store in a cool, dry place.**

Manufactured  
by:



148 Longmeadow Road  
Taunton, MA 02780  
(508) 824-7292

**WARNING:** Do not feed to sheep or goats producing milk for food.

**CAUTION:** Do not use in feeds containing bentonite.

**CAUTION:** THIS FEED CONTAINS ADDED COPPER DO NOT FEED TO SHEEP OR OTHER COPPER SENSITIVE SPECIES

Feed at a rate of 22.7 mg decoquinatate per 100lb of body weight (0.5 mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

### Feeding Guide:

Body Weight	Lbs. of feed (to provide 22.7 mg Decoquinatate/100 lbs bodyweight)
25 lbs	0.50
50 lbs	1.00
75 lbs	1.50
100 lbs	2.00

*The suggested feeding program is for use as a guide only. The animal's requirements may change due to breed, environment and management.*

**Always Provide plenty of clean fresh water.**