



# EVERYDAY Turkey Starter Crumbles

## MEDICATED Type C

**ACTIVE DRUG INGREDIENT: MONENSIN 72g/ton**

For the prevention of coccidiosis caused by Eimeria adenoeides, E. melagrimitis and E. gallopavonis.

### GUARANTEED ANALYSIS

CRUDE PROTEIN		(MIN)	28.0%
LYSINE		(MIN)	1.60%
METHIONINE		(MIN)	0.50%
CRUDE FAT		(MIN)	4.50%
CRUDE FIBER		(MAX)	4.00%
CALCIUM	(MIN)	1.30%	(MAX) 1.50%
PHOSPHORUS		(MIN)	0.70%
SALT	(MIN)	0.06%	(MAX) 0.08%

### INGREDIENTS

Ground Yellow Corn, Soybean Meal, Fish Meal, Corn Distillers Dried Grains w/Solubles, Alfalfa Meal, Fat-Vegetable Blend, Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Salt, DL-Methionine, L-Lysine, Processed Grain By-Products, Manganous Oxide, Zinc Oxide, Vitamin E Supplement, Ferrous Sulfate, Niacin Supplement, Roughage Products, Copper Sulfate, Calcium Pantothenate, Riboflavin Supplement, Mineral Oil, Menadione Nicotinamide Bisulfite, Vitamin A Supplement, Sodium Selenite, Pyridoxine Hydrochloride, Vitamin D3 Supplement, Thiamine Mononitrate, Folic Acid, Vitamin B12 Supplement, Biotin, Ethylenediamine Dihydriodide, Natural Flavors, Choline Chloride, Stresseze

**Ingredients as defined by AAFCO.**

**Do not store on ground, store in a cool, dry place.**

Manufactured  
by:



148 Longmeadow Road  
Taunton, MA 02780  
(508) 824-7292

**DIRECTIONS FOR USE:** Feed continuously to growing turkeys only as the sole ration. The optimum level depends on the severity of coccidiosis exposure.

**WARNING:** Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age.

**CAUTION:** Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis. Not for replacement chickens intended to become broiler breeding chickens.

*The suggested feeding program is for use as a guide only. The animal's requirements may change due to breed, environment and management.*